


IN THE CLAIMS:

1. (Currently amended) A process for treatment of at least one condition chosen from seborrhoea of the skin and scalp, disorders associated with seborrhoea, and disorders associated with microorganisms of the genus *Propionibacterium*, said process comprising:

applying to an area in need of said treatment at least one compound chosen from polyamino acid derivatives of formula (I) and salts thereof,




in which:

X is chosen from O, S, NH and NR" wherein R" is chosen from saturated and unsaturated, linear and branched C₁₋₆ hydrocarbon-based radicals;

R₁ is chosen from:

- (i) hydrogen;
- (ii) linear and branched, saturated and unsaturated C₁₋₄₀ hydrocarbon-based radicals,
- (iii) radicals of the formula




FINNEGAN
HENDERSON
FARABOW
GARRETT &
DUNNER LLP

1300 I Street, NW
Washington, DC 20005
202.408.4000
Fax 202.408.4400
www.finnegan.com


wherein s is a number chosen from 0, 1, 2, 3 and 4; and R₄ is chosen from hydrogen and radicals chosen from -NH₂, -OH, -SH, -CHOHCH₃, -CONH₂, -NH-C(NH₂)=NH, -C₆H₅,

-C₆H₄OH and



and;

(iv) radicals of the formula



wherein m is a number chosen from 3, 4 and 5;

R₂ is chosen from hydrogen; saturated and unsaturated, linear and branched C₁₋₈ hydrocarbon-based radicals; and radicals chosen from -CH₂C₆H₅, -CH₂C₆H₄OH, -CH₂OH,

-CHOHCH₃, -(CH₂)_t-NH₂, wherein t is a number chosen from 3, 4 and 5;

R₃ is chosen from hydrogen and saturated and unsaturated, linear and branched C₁₋₆ hydrocarbon-based radicals; and

n is a number greater than 1 chosen such that the number average molecular weight of the polyamino acid derivative ranges from ~~400~~ 150 to 200 000;

wherein the repeating unit may be identical or different for the same derivative.

FINNEGAN
HENDERSON
FARABOW
GARRETT &
DUNNER LLP

1300 I Street, NW
Washington, DC 20005
202.408.4000
Fax 202.408.4400
www.finnegan.com

2. (Original) A process according to claim 1, wherein said microorganisms are *Propionibacterium acnes*.

3. (Original) A process according to claim 1, wherein said microorganisms are *Propionibacterium granulosum*.

4. (Original) A process according to claim 1, wherein R_1 is chosen from linear and branched, saturated and unsaturated C_{1-40} hydrocarbon-based radicals substituted with at least one hydroxyl radical, at least one radical -NRR', or at least one hydroxyl radical and at least one radical -NRR', wherein R and R', which may be identical or different, are chosen from hydrogen and saturated and unsaturated, linear and branched C_{1-6} hydrocarbon-based radicals.

5. (Original) A process according to claim 1, wherein R_1 is chosen from linear and branched, saturated and unsaturated C_{1-40} hydrocarbon-based radicals interrupted with at least one hetero atom chosen from N, O and Si.

6. (Original) A process according to claim 1, wherein said at least one compound is administered in the form of a cosmetic composition.

7. (Original) A process according to claim 6, wherein the treatment comprises the cosmetic treatment of at least one disorder chosen from seborrhoeic dermatitis, acne, greasy skin with a tendency towards acne, and hyperseborrhoea.

FINNEGAN
HENDERSON
FARABOW
GARRETT &
DUNNER LLP

1300 I Street, NW
Washington, DC 20005
202.408.4000
Fax 202.408.4400
www.finnegan.com

8. (Original) A process according to claim 1, wherein said at least one compound is administered in the form of a pharmaceutical composition.

9. (Original) A process according to claim 8, in which the pharmaceutical composition is administered for treating at least one disorder chosen from seborrhoeic dermatitis, acne, greasy skin with a tendency towards acne and hyperseborrhoea.

10. (Original) A process according to claim 1, wherein in said polyamino acid derivatives of formula (I) and salts thereof, at least one of the following definitions apply:

X is chosen from O, S, NH and NR", wherein R" is chosen from saturated and unsaturated, linear and branched C₁₋₆ hydrocarbon-based radicals;

R₁ is chosen from linear and branched, saturated and unsaturated C₈₋₄₀ hydrocarbon-based radicals,

R₂ is hydrogen;

R₃ is chosen from saturated, linear and branched C₁₋₆ hydrocarbon-based radicals; and

n is chosen from a number ranging from 2 to 100 and a number chosen such that the number average molecular weight of said polyamino acid derivative ranges from 150 to 10,000.

11. (Original) A process according to claim 10, wherein R₃ is chosen from methyl and ethyl radicals.

FINNEGAN
HENDERSON
FARABOW
GARRETT &
DUNNER LLP

1300 I Street, NW
Washington, DC 20005
202.408.4000
Fax 202.408.4400
www.finnegan.com

12. (Original) A process according to claim 10, wherein R_1 is chosen from linear and branched, saturated and unsaturated C_{8-40} hydrocarbon-based radicals substituted with at least one hydroxyl radical, at least one radical $-NRR'$, or at least one hydroxyl radical and at least one radical $-NRR'$, wherein R and R' , which may be identical or different, are chosen from hydrogen and saturated and unsaturated, linear and branched C_{1-6} hydrocarbon-based radicals.

13. (Original) A process according to claim 10, wherein R_1 is chosen from linear and branched, saturated and unsaturated C_{8-40} hydrocarbon-based radicals interrupted with at least one hetero atom chosen from N, O and Si.

14. (Currently amended) A process according to claim 44 13, wherein n is chosen from a number ranging from 2 to 100.

15. (Original) A process according to claim 14, wherein n is a number chosen such that the number average molecular weight of said polyamino acid derivative ranges from 150 to 10,000.

16. (Original) A process according to claim 10, wherein:
X is chosen from O, S, NH and NR'' , wherein R'' is chosen from saturated and unsaturated, linear and branched C_{1-6} hydrocarbon-based radicals;

R_1 is chosen from linear and branched, saturated and unsaturated C_{8-40} hydrocarbon-based radicals,

FINNEGAN
HENDERSON
FARABOW
GARRETT &
DUNNER LLP

1300 I Street, NW
Washington, DC 20005
202.408.4000
Fax 202.408.4400
www.finnegan.com

R₂ is hydrogen;

R₃ is chosen from saturated, linear and branched C₁₋₆ hydrocarbon-based radicals; and

n is chosen from a number ranging from 2 to 100 and a number chosen such that the number average molecular weight of said polyamino acid derivative ranges from 150 to 10,000.

17. (Original) A process according to claim 1, wherein in said polyamino acid derivatives of formula (I) and salts thereof, at least one of the following definitions apply:

X is chosen from O, S and NH;

R₁ is chosen from linear and branched, saturated C₁₀₋₂₄ hydrocarbon-based radicals; and linear and branched unsaturated hydrocarbon-based radicals;

R₂ is hydrogen;

R₃ is a methyl radical; and

n is chosen from a number ranging from 4 to 50 and a number chosen such that the number average molecular weight of said polyamino acid derivative ranges from 300 to 8,000.

18. (Original) A process according to claim 17, wherein n is chosen from a number ranging from 4 to 50.

FINNEGAN
HENDERSON
FARABOW
GARRETT &
DUNNER LLP

1300 I Street, NW
Washington, DC 20005
202.408.4000
Fax 202.408.4400
www.finnegan.com

19. (Original) A process according to claim 17, wherein n is a number chosen such that the number average molecular weight of said polyamino acid derivative ranges from 300 to 8,000.

20. (Original) A process according to claim 17, wherein X is NH.

21. (Original) A process according to claim 17, wherein R₁ is chosen from linear and branched, saturated C₁₀₋₂₄ hydrocarbon-based radicals substituted with at least one hydroxyl radical.

22. (Original) A process according to claim 21, wherein said linear and branched, saturated C₁₀₋₂₄ hydrocarbon-based radicals are substituted with 1, 2, 3, or 4 hydroxyl radicals.

23. (Original) A process according to claim 17, wherein R₁ is chosen from linear and branched unsaturated hydrocarbon-based radicals substituted with at least one hydroxyl radical.

24. (Original) A process according to claim 1, wherein:

X is chosen from O, S and NH;

R₁ is chosen from linear and branched, saturated C₁₀₋₂₄ hydrocarbon-based radicals; and linear and branched unsaturated hydrocarbon-based radicals;

R₂ is hydrogen;

AL
amt

FINNEGAN
HENDERSON
FARABOW
GARRETT &
DUNNER LLP

1300 I Street, NW
Washington, DC 20005
202.408.4000
Fax 202.408.4400
www.finnegan.com

R_3 is a methyl radical; and

n is chosen from a number ranging from 4 to 50 and a number chosen such that the number average molecular weight of said polyamino acid derivative ranges from 300 to 8,000.

25. (Original) A process according to claim 1, wherein said at least one compound is present in said composition in an amount ranging from 0.001% to 30% by weight, relative to the total weight of the composition.

26. (Original) A process according to claim 25, wherein said at least one compound is present in said composition in an amount ranging from 0.01% to 15% by weight, relative to the total weight of the composition.

27. (Original) A process according to claim 26, wherein said at least one compound is present in said composition in an amount ranging from 0.5% to 5% by weight, relative to the total weight of the composition.

28. (Original) A process according to claim 10, wherein said at least one compound is applied in the form of a composition chosen from a cosmetic composition and a pharmaceutical composition.

PH
CMT

FINNEGAN
HENDERSON
FARABOW
GARRETT &
DUNNER LLP


1300 I Street, NW
Washington, DC 20005
202.408.4000
Fax 202.408.4400
www.finnegan.com

29. (Original) A process according to claim 17, wherein said at least one compound is applied in the form of a composition chosen from a cosmetic composition and a pharmaceutical composition.

30. (Original) A process according to claim 1, wherein said at least one compound is applied to at least one area chosen from the skin and the scalp.

31. (Withdrawn) A process for the manufacture of a composition for treatment of at least one condition chosen from seborrhoea of the skin and scalp, disorders associated with seborrhoea, and disorders associated with microorganisms of the genus *Propionibacterium*, said process comprising:

including in said composition at least one poly amino acid derivative chosen from formula (I) and salts thereof ,



in which:

X is chosen from O, S, NH and NR" with R" is chosen from saturated and unsaturated, linear and branched C₁₋₆ hydrocarbon-based radicals;


R₁ is chosen from:

- (i) hydrogen;
- (ii) linear and branched, saturated and unsaturated C₁₋₄₀ hydrocarbon-based radicals,


FINNEGAN
HENDERSON
FARABOW
GARRETT &
DUNNER LLP

1300 I Street, NW
Washington, DC 20005
202.408.4000
Fax 202.408.4400
www.finnegan.com

(iii) radicals of the formula




wherein s is a number chosen from 0, 1, 2, 3 and 4; and R₄ is chosen from hydrogen and radicals chosen from -NH₂, -OH, -SH, -CHOHCH₃, -CONH₂, -NH-C(NH₂)=NH, -C₆H₅, -C₆H₄OH and



and;

(iv) radicals of the formula



wherein m is a number chosen from 3, 4 and 5;

- R₂ is chosen from hydrogen; saturated and unsaturated, linear and branched C₁₋₈ hydrocarbon-based radicals; and radicals chosen from -CH₂C₆H₅, -CH₂C₆H₄OH, -CH₂OH, -CHOHCH₃, -(CH₂)_t-NH₂ wherein t is a number chosen from 3, 4 and 5;

R₃ is chosen from hydrogen and saturated and unsaturated, linear and branched C₁₋₆ hydrocarbon-based radicals; and

FINNEGAN
HENDERSON
FARABOW
GARRETT &
DUNNER LLP

1300 I Street, NW
Washington, DC 20005
202.408.4000
Fax 202.408.4400
www.finnegan.com

n is a number greater than 1 chosen such that the number average molecular weight of the polyamino acid derivative ranges from 100 to 200 000;
wherein the repeating unit may be identical or different for the same derivative.

32. (Withdrawn) A process according to claim 31, wherein said microorganisms are *Propionibacterium acnes*.

33. (Withdrawn) A process according to claim 31, wherein said microorganisms are *Propionibacterium granulosum*.

34. (Withdrawn) A process according to claim 31, wherein R₁ is chosen from linear and branched, saturated and unsaturated C₁₋₄₀ hydrocarbon-based radicals substituted with at least one hydroxyl radical, at least one radical -NRR', or at least one hydroxyl radical and at least one radical -NRR', wherein R and R', which may be identical or different, are chosen from hydrogen and saturated and unsaturated, linear and branched C₁₋₆ hydrocarbon-based radicals.

35. (Withdrawn) A process according to claim 31, wherein R₁ is chosen from linear and branched, saturated and unsaturated C₁₋₄₀ hydrocarbon-based radicals interrupted with at least one hetero atom chosen from N, O and Si.

FINNEGAN
HENDERSON
FARABOW
GARRETT &
DUNNER LLP

1300 I Street, NW
Washington, DC 20005
202.408.4000
Fax 202.408.4400
www.finnegan.com

36. (Withdrawn) A process according to claim 31, wherein in said polyamino acid derivatives of formula (I) and salts thereof, at least one of the following definitions apply:

X is chosen from O, S, NH and NR", wherein R" is chosen from saturated and unsaturated, linear and branched C₁₋₆ hydrocarbon-based radicals;

R₁ is chosen from linear and branched, saturated and unsaturated C₈₋₄₀ hydrocarbon-based radicals,

R₂ is hydrogen;

R₃ is chosen from saturated, linear and branched C₁₋₆ hydrocarbon-based radicals; and

n is chosen from a number ranging from 2 to 100 and a number chosen such that the number average molecular weight of said polyamino acid derivative ranges from 150 to 10,000.

37. (Withdrawn) A process according to claim 36, wherein R₃ is chosen from methyl and ethyl radicals.

38. (Withdrawn) A process according to claim 36, wherein R₁ is chosen from linear and branched, saturated and unsaturated C₈₋₄₀ hydrocarbon-based radicals substituted with at least one hydroxyl radical, at least one radical -NRR', or at least one hydroxyl radical and at least one radical -NRR', wherein R and R', which may be identical or different, are chosen from hydrogen and saturated and unsaturated, linear and branched C₁₋₆ hydrocarbon-based radicals.

FINNEGAN
HENDERSON
FARABOW
GARRETT &
DUNNER LLP

1300 I Street, NW
Washington, DC 20005
202.408.4000
Fax 202.408.4400
www.finnegan.com

39. (Withdrawn) A process according to claim 36, wherein R_1 is chosen from linear and branched, saturated and unsaturated C_{8-40} hydrocarbon-based radicals interrupted with at least one hetero atom chosen from N, O and Si.

40. (Withdrawn) A process according to claim 36, wherein n is chosen from a number ranging from 2 to 100.

41. (Withdrawn) A process according to claim 36, wherein n is a number chosen such that the number average molecular weight of said polyamino acid derivative ranges from 150 to 10,000.

42. (Withdrawn) A process according to claim 36, wherein:

X is chosen from O, S, NH and NR'' , wherein R'' is chosen from saturated and unsaturated, linear and branched C_{1-6} hydrocarbon-based radicals;

R_1 is chosen from linear and branched, saturated and unsaturated C_{8-40} hydrocarbon-based radicals,

R_2 is hydrogen;

R_3 is chosen from saturated, linear and branched C_{1-6} hydrocarbon-based radicals; and

n is chosen from a number ranging from 2 to 100 and a number chosen such that the number average molecular weight of said polyamino acid derivative ranges from 150 to 10,000.

FINNEGAN
HENDERSON
FARABOW
GARRETT &
DUNNER LLP

1300 I Street, NW
Washington, DC 20005
202.408.4000
Fax 202.408.4400
www.finnegan.com

43. (Withdrawn) A process according to claim 31, wherein in said polyamino acid derivatives of formula (I) and salts thereof, at least one of the following definitions apply:

X is chosen from O, S and NH;

R₁ is chosen from linear and branched, saturated C₁₀₋₂₄ hydrocarbon-based radicals; and linear and branched unsaturated hydrocarbon-based radicals;

R₂ is hydrogen;

R₃ is a methyl radical; and

n is chosen from a number ranging from 4 to 50 and a number chosen such that the number average molecular weight of said polyamino acid derivative ranges from 300 to 8,000.

44. (Withdrawn) A process according to claim 43, wherein n is chosen from a number ranging from 4 to 50.

45. (Withdrawn) A process according to claim 43, wherein n is a number chosen such that the number average molecular weight of said polyamino acid derivative ranges from 300 to 8,000.

46. (Withdrawn) A process according to claim 43, wherein X is NH.

FINNEGAN
HENDERSON
FARABOW
GARRETT &
DUNNER ^{LLP}

1300 I Street, NW
Washington, DC 20005
202.408.4000
Fax 202.408.4400
www.finnegan.com

47. (Withdrawn) A process according to claim 43, wherein R_1 is chosen from linear and branched, saturated C_{10-24} hydrocarbon-based radicals substituted with at least one hydroxyl radical.

48. (Withdrawn) A process according to claim 47, wherein said linear and branched, saturated C_{10-24} hydrocarbon-based radicals are substituted with 1, 2, 3, or 4 hydroxyl radicals.

49. (Withdrawn) A process according to claim 43, wherein R_1 is chosen from linear and branched unsaturated hydrocarbon-based radicals substituted with at least one hydroxyl radical.

50. (Withdrawn) A process according to claim 31, wherein said at least one polyamino acid derivative is present in said composition in an amount ranging from 0.001% to 30% by weight, relative to the total weight of the composition.

51. (Withdrawn) A process according to claim 50, wherein said at least one polyamino acid derivative is present in said composition in an amount ranging from 0.01% to 15% by weight, relative to the total weight of the composition.


52. (Withdrawn) A process according to claim 51, wherein said at least one polyamino acid derivative is present in said composition in an amount ranging from 0.5% to 5% by weight, relative to the total weight of the composition.

FINNEGAN
HENDERSON
FARABOW
GARRETT &
DUNNER LLP

1300 I Street, NW
Washington, DC 20005
202.408.4000
Fax 202.408.4400
www.finnegan.com

53. (Withdrawn) A process according to claim 31, wherein said composition is a pharmaceutical composition.

54. (Withdrawn) An anti-seborrhoeic composition comprising,
a physiologically acceptable medium; and
an effective amount of at least one polyamino acid derivative of formula (I) and salts thereof,




in which:

X is chosen from O, S, NH and NR" wherein R" is chosen from saturated and unsaturated, linear and branched C₁₋₆ hydrocarbon-based radicals;

R₁ is chosen from:


- (i) hydrogen;
- (ii) linear and branched, saturated and unsaturated C₁₋₄₀ hydrocarbon-based radicals,
- (iii) radicals of the formula



FINNEGAN
HENDERSON
FARABOW
GARRETT &
DUNNER LLP


1300 I Street, NW
Washington, DC 20005
202.408.4000
Fax 202.408.4400
www.finnegan.com

wherein s is a number chosen from 0, 1, 2, 3 and 4; and R₄ is chosen from hydrogen and radicals chosen from -NH₂, -OH, -SH, -CHOHCH₃, -CONH₂, -NH-C(NH₂)=NH, -C₆H₅, -C₆H₄OH and



and;

(iv) radicals of the formula



wherein m is a number chosen from 3, 4 and 5;

R₂ is chosen from hydrogen; saturated and unsaturated, linear and branched C₁₋₈ hydrocarbon-based radicals; and radicals chosen from -CH₂C₆H₅, -CH₂C₆H₄OH, -CH₂OH, -CHOHCH₃, -(CH₂)_t-NH₂ wherein t is a number chosen from 3, 4 and 5;

R₃ is chosen from hydrogen and saturated and unsaturated, linear and branched C₁₋₆ hydrocarbon-based radicals; and


n is a number greater than 1 chosen such that the number average molecular weight of the polyamino acid derivative ranges from 100 to 200 000;

wherein the repeating unit may be identical or different for the same derivative.

55. (Withdrawn) An anti-seborrhoeic composition according to claim 54,

wherein said composition is an anti-acne composition.

56. (Withdrawn) An anti-bacterial composition comprising,
a physiologically acceptable medium; and
an effective amount of at least one polyamino acid derivative of formula (I) and
salts thereof for treating bacteria,




in which:

X is chosen from O, S, NH and NR" wherein R" is chosen from saturated and
unsaturated, linear and branched C₁₋₆ hydrocarbon-based radicals;

R₁ is chosen from:


- (i) hydrogen;
- (ii) linear and branched, saturated and unsaturated C₁₋₄₀ hydrocarbon-based
radicals,
- (iii) radicals of the formula



wherein s is a number chosen from 0, 1, 2, 3 and 4; and R₄ is chosen from hydrogen
and radicals chosen from -NH₂, -OH, -SH, -CHOHCH₃, -CONH₂, -NH-C(NH₂)=NH, -
C₆H₅, -C₆H₄OH and


FINNEGAN
HENDERSON
FARABOW
GARRETT &
DUNNER LLP

1300 I Street, NW
Washington, DC 20005
202.408.4000
Fax 202.408.4400
www.finnegan.com



and;

(iv) radicals of the formula



wherein m is a number chosen from 3, 4 and 5;

R₂ is chosen from hydrogen; saturated and unsaturated, linear and branched C₁₋₈ hydrocarbon-based radicals; and radicals chosen from -CH₂C₆H₅, -CH₂C₆H₄OH, -CH₂OH, -CHOHCH₃, -(CH₂)_t-NH₂ wherein t is a number chosen from 3, 4 and 5;

R₃ is chosen from hydrogen and saturated and unsaturated, linear and branched C₁₋₆ hydrocarbon-based radicals; and

n is a number greater than 1 chosen such that the number average molecular weight of the polyamino acid derivative ranges from 100 to 200 000;


wherein the repeating unit may be identical or different for the same derivative.

57. (Withdrawn) An anti-bacterial composition according to claim 56, wherein said composition is an anti-acne composition.

58. (Withdrawn) An antibacterial composition according to claim 55, wherein the bacteria is of the genus *Propionibacterium*.

FINNEGAN
HENDERSON
FARABOW
GARRETT &
DUNNER LLP

1300 I Street, NW
Washington, DC 20005
202.408.4000
Fax 202.408.4400
www.finnegan.com



59. (Withdrawn) An antibacterial composition according to claim 58, wherein the bacteria is at least one of *Propionibacterium acnes* and *Propionibacterium granulosum*.

FINNEGAN
HENDERSON
FARABOW
GARRETT &
DUNNER ^{LLP}

1300 I Street, NW
Washington, DC 20005
202.408.4000
Fax 202.408.4400
www.finnegan.com